

## NOTES ON GEOGRAPHIC DISTRIBUTION

### Reptilia, Squamata, Leiosauridae, *Urostrophus vautieri*: Distribution extension and geographic distribution map

Rodrigo Costa Santos<sup>1</sup>  
Priscila Silva Lucas<sup>1</sup>  
Bernadete Maria de Sousa<sup>2</sup>  
Iara Alves Novelli<sup>1,2</sup>

<sup>1</sup> Centro Universitário de Lavras - UNILAVRAS, Laboratório de Zoologia.  
Rua Padre José Poggel, 506, Centenário. CEP 37200-000. Lavras, MG, Brazil.

<sup>2</sup> Universidade Federal de Juiz de Fora, Instituto de Ciências Biológicas, Laboratório de Herpetologia.  
Campus Universitário, Bairro Martelos, CEP 36036-330. Juiz de Fora, MG, Brazil.  
E-mail: iaranovelli27@gmail.com

The genus *Urostrophus* comprises two recognized species, *Urostrophus vautieri* Duméril & Bibron, 1837 and *Urostrophus gallardoi* Etheridge & Williams, 1991, both from southern South America (Etheridge and Williams 1991).

*Urostrophus vautieri* is typically found in the Brazilian Atlantic rainforest biome (Etheridge and Williams 1991; Pellegrino et al. 1999) in the states of southeastern (Espírito Santo, Minas Gerais, Rio de Janeiro, and São Paulo) and southern regions (Santa Catarina, Paraná, and Rio Grande do Sul) (Figure 1) (Von Ihering 1898; Lema 1958; 1994; 2002; Ruschi 1966; Etheridge and Williams 1991; Sazima and Haddad 1992; Pellegrino et al. 1999; Bérnills et al. 2001; 2007; Rocha et al. 2004; Ribeiro and Sousa 2006; Kunz et al. 2007).

On 26 September 2008 and 21 November 2008, we collect two adult males of *Urostrophus vautieri* (Figure 2 and 3) in *Reserva do Boqueirão* (21°20'47" S, 44°59'27" W, 1250 m elevation), in riparian forest included in the *Cerrado* biome, in municipality of Ingaí, state of Minas Gerais. Voucher specimens (CRLZ 000065, 000071) are deposited at the *Coleção de Répteis do Laboratório de Zoologia, Centro Universitário de Lavras - UNILAVRAS*, municipality of Lavras,

state of Minas Gerais, Brazil. Collecting permits was authorized by Brazilian Institute of Environment and Renewable Natural Resources (*Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis - IBAMA*) (Process number - SISBIO/14740-1).



**Figure 1.** Distribution of the lizard *Urostrophus vautieri* in Brazil .

NOTES ON GEOGRAPHIC DISTRIBUTION



**Figure 2.** Adult male *U. vautieri* (CRLZ 000065) collected at *Reserva do Boqueirão*, municipality of Ingai, state of Minas Gerais. Photo by Rodrigo C. Santos.



**Figure 3.** Adult male *U. vautieri* (CRLZ 000065) collected at *Reserva do Boqueirão*, municipality of Ingai, state of Minas Gerais. Photo by Rodrigo C. Santos.


## NOTES ON GEOGRAPHIC DISTRIBUTION

This species was registered in the state of Minas Gerais (Figure 4) in Lagoa Santa (19°38' S, 43°52' W) (Etheridge and Williams 1991), Engenheiro Trompowski (21°18' S, 46°17' W) (Etheridge and Williams 1991), Antonio Carlos (21°19' S, 43°45' W) (Etheridge and Williams 1991), Poços de Caldas (21°48' S, 46°34' W) (Etheridge and Williams 1991), Santa Rita da Extrema (22°52' S, 46°19' W) (Etheridge and Williams 1991), and Juiz de Fora (21°46' S, 43°22' W) (Ribeiro and Sousa 2006). In other states (Figure 4), *U. vautieri* was registered close to the state of Minas Gerais, in Campos do Jordão (22°45' S, 45°34' W) and Serra Negra (22°37' S, 46°42' W), state of São Paulo (Etheridge and Williams 1991), and in municipality of Itatiaia (22°23' S, 44°39' W),

state of Rio de Janeiro (Etheridge and Williams 1991).

The records here reported extend the known geographical distribution of *U. vautieri* ca. 129 km west from the nearest record in the state of Minas Gerais, in municipality of Antônio Carlos, and ca. 120 km northwest from the nearest record in the state of Rio de Janeiro, in municipality of Itatiaia (Figure 4).

Some authors characterize this species as typical of the Atlantic rainforest biome (Etheridge and Williams 1991, Pellegrino et al. 1999). However, the specimens of the current study were collected in a Biological Reserve included in *Cerrado*. This is the first record of *U. vautieri* in this biome.



**Figure 4.** Distribution of the lizard *Urostrophus vautieri* in Minas Gerais, São Paulo, and Rio de Janeiro states. The red circle represents the municipality of Ingai (new distribution record), state of Minas Gerais. Dotted arrow indicates the municipality of Itatiaia, state of Rio de Janeiro. The black arrow indicates the municipality of Antonio Carlos, state of Minas Gerais.

### Acknowledgements

We thank to Prof. Fernando Frieiro, Fabiano B. Silva, and Fabiano M. Vieira for logistical support. RCS was supported by a student fellowship from PROBIC/FAPEMIG/UNILAVRAS.

## NOTES ON GEOGRAPHIC DISTRIBUTION

### Literature Cited

- Bérnills, R. S., A. R. Giraudo, S. Carreira and S. Z. Cechin. 2007. Répteis das porções subtropical e temperada da Região Neotropical. *Ciência & Ambiente* 35: 101-136.
- Bérnills, R. S., M. A. Batista and P. W. Bertelli. 2001. Cobras e lagartos do Vale: levantamento das espécies de Squamata (Reptilia, Lepidosauria) da Bacia do Rio Itajai, Santa Catarina, Brasil. *Revista de estudos ambientais* 3: 69-79.
- Etheridge, R. and E. E. Williams. 1991. A review of the South American lizard genera *Urostrophus* and *Anisolepis* (Squamata: Iguania: Polychridae). *Bulletin of the Museum of Comparative Zoology* 152: 317-361.
- Kunz, T. S., I. R. Ghizoni-Jr., W. L. A. Santos and P. A. Hartmann. 2007. Nota sobre a coleção herpetológica da Universidade Federal de Santa Catarina (UFSC). *Biotemas* 20(3): 127-132.
- Lema, T. 1958. Notas sobre os répteis do estado do Rio Grande do Sul. 1. Introdução ao estudo dos répteis do Estado do Rio Grande do Sul. 1. Histórico. *Iheringia (Zoologia)* 10: 5-18.
- Lema, T. 1994. Lista comentada dos répteis ocorrentes no Rio Grande do Sul, Brasil. *Comunicações do Museu de Ciências da PUC do Rio Grande do Sul, Série Zoologia*, 7: 41-150.
- Lema, T. 2002. Os répteis do Rio Grande do Sul. Atuais e fósseis – Biogeografia – Ofidismo. Porto Alegre: EDIPUCRS. 264 p.
- Pellegrino, K. C., C. E. V. Bertolotto, M. T. Rodrigues and Y. Yonenaga-Yassuda. 1999. Banding patterns, heteromorphic sex chromosomes and Ag-stained NORs after pachytene stages in the meiosis of the Brazilian lizard *Urostrophus vautieri* (Squamata, Polychrotidae). *Caryologia* 52(1-2): 21-26.
- Ribeiro, L. B. and B. M. Sousa. 2006. *Urostrophus vautieri* (NCN). Skin shedding. *Herpetological Review* 37(3): 348.
- Rocha, C. F. D., H. G. Bergallo, J. P. Pombal Jr., L. Geise, M. Van Sluys, R. Fernandes and U. Caramaschi. 2004. Lista de anfíbios, répteis e mamíferos do Estado do Rio de Janeiro, Sudeste do Brasil. *Publicações Avulsas do Museu Nacional* 104: 1-24.
- Ruschi, A. 1966. Lista do répteis do estado do Espírito Santo. *Boletim do Museu de Biologia Prof. Mello-Leitão, Zoologia*, 26A: 1-6.
- Sazima, I. and C. F. B. Haddad. 1992. Répteis da Serra do Japi: notas sobre História Natural; p. 212-236 In: L. P. C. Morellato (ed.). *História Natural da Serra do Japi: ecologia e preservação de uma área florestal do sudeste do Brasil*. Campinas: Editora da UNICAMP/FAPESP.
- Von Ihering, H. 1898. Contributions to the herpetology of Sao Paulo, Brazil: I. Proceedings of the Academy of Natural Sciences of Philadelphia 50: 101-109.

Received March 2009

Accepted July 2009

Published online September 2009